

ACEPROMAZINE 1 mg/mL		Dosage (IM or SQ) 0.05 mg/kg, capped at 1 mg at 31 ≥ lbs (14 ≥ kg)
Weight (Kgs)		mL
1.0 - 2.5		0.1
3.0 - 4.5		0.2
5.0 - 9.0		0.5
9.5 - 13.5		0.7
		Weight (Lbs)
		2.2 - 5.5
		6.6 - 10.0
		11.0 - 19.8
		21.0 - 29.7

ACEPROMAZINE 10 mg/mL		Dosage (IM or SQ) 0.05 mg/kg, capped at 1 mg at 31 ≥ lbs (14 ≥ kg)
Weight (Kgs)		mL
14.0 ≥		0.1
		Weight (Lbs)
		30.8 ≥

HYDROMORPHONE 10 mg/mL		Dosage (IM or SQ) 0.1 mg/kg
Weight (Kgs)		mL
1 - 2.5		0.02
3 - 4.5		0.04
5 - 7		0.06
7.5 - 9		0.08
9.5 - 11.5		0.10
12 - 13.5		0.13
14 - 16		0.15
16.5 - 18		0.17
18.5 - 20.5		0.20
21 - 23		0.22
23.5 - 25		0.24
25.5 - 27		0.26
27.5 - 29.5		0.28
30 - 32		0.30
32.5 - 34		0.32
34.5 - 36.5		0.34
37 - 38.5		0.38
39 - 41		0.40
41.5 - 43		0.42
43.5 - 45.5		0.44
46 - 48		0.46
48.5 - 50		0.48
		Weight (Lbs)
		2.0 - 5.5
		6.0 - 9.9
		11.0 - 15.0
		16.5 - 20.0
		20.9 - 25.3
		26.4 - 29.7
		30.8 - 35.0
		36.3 - 40.0
		40.7 - 45.1
		46.0 - 50.0
		51.7 - 55.0
		56.1 - 60.0
		60.5 - 64.9
		66.0 - 70.4
		71.5 - 75.0
		75.9 - 80.3
		81.0 - 84.7
		86.0 - 90.0
		91.3 - 95.0
		95.7 - 100.1
		101.0 - 105.0
		106.7 - 110.0

Consider giving premed intramuscularly (IM) vs. subcutaneously (SQ) to increase intensity and onset of analgesia and sedation.