

Your pet was found to be nauseous while staying with us. Nausea is a common occurrence after surgery involving anesthesia, and usually resolves on its own. Please feed smaller amounts of your pet's normal food for a day or so until their system returns to normal. We have given your pet one or more of the following treatments to help with nausea and hydration:

•	mL Cerenia (anti-nausea)	
• _	mL Famotidine (anti-acid)	
•	mL subcutaneous fluids (for hydration)	
	niting continues longer than 12-24 hours, or if you 1079 (local office hours) to schedule a recheck, or	
ASI	PCA Spay/Neuter Alliance	Client Nausea Handout
after s	pet was found to be nauseous while staying with surgery involving anesthesia, and usually resolves or pet's normal food for a day or so until their systoet one or more of the following treatments to hel	on its own. Please feed smaller amounts em returns to normal. We have given
•	mL Cerenia (anti-nausea)	
•	mL Famotidine (anti-acid)	
•	mL subcutaneous fluids (for hydration)	
	niting continues longer than 12-24 hours, or if you 1079 (local office hours) to schedule a recheck, or	
ASP	Spay/Neuter Alliance	Client Nausea Handout
after s	pet was found to be nauseous while staying with surgery involving anesthesia, and usually resolves or pet's normal food for a day or so until their system one or more of the following treatments to hele	on its own. Please feed smaller amounts em returns to normal. We have given
•	mL Cerenia (anti-nausea)	
•	mL Famotidine (anti-acid)	
•	mL subcutaneous fluids (for hydration)	

If vomiting continues longer than 12-24 hours, or if you have any questions, please call **(828) 252-2079** (local office hours) to schedule a recheck, or **(855) 434-9285** (after-hours helpline).